

Dietary Formulation and Nutritional Composition of Cereal Based Complementary Food

Adefisola Bola Adepaju¹ and Samson Ishola Ibiroko^{2*}

¹Department of Food Science and Technology, College of Agriculture, Joseph Ayo Babalola University, Ikeji- Arakeji, Nigeria

²Department of Food Science and Technology, Faculty of Technology, Obafemi Awolowo University Ile-Ife, Osun State, Nigeria

***Corresponding Author:** Samson Ishola Ibiroko, Department of Food Science and Technology, Faculty of Technology, Obafemi Awolowo University Ile-Ife, Osun State, Nigeria.

Received: April 23, 2019; **Published:** July 08, 2019

Abstract

Globally study revealed that, Nigeria loses about 2,300 under-five year olds and 145 women of child bearing age, making the country the second largest contributor to under-five and maternal mortality rate in the world. The purpose of the study is to formulate and determine of Nutritional composition of Cereal based complementary food that will have adequate nutrient to prevent children and infants mortality. The Cereal based complementary food composed of Sorghum 70% Soybean 30% (SS), Millet 70% Soybean 30% (MS), Sorghum 35% Millet 35%, Soybean 30% (SMS); Basal Dietary (BD); and Commercial Food (CF). The result revealed that that bioassay of the experimental animal Bioassay BV % ranged from 66 - 75% NPU% 64 - 72 PER 1.2 - 4.2 and NPR 1.7 - 3.3 were compared favourably with commercial samples. Chemical composition in the sample dietary met the recommended daily requirement. In conclusion raw material to produce cereal based complementary food is available for baby and infant in most part of the developing countries. Millet grain containing dietary promoted the growth response next to commercial diet. It contains alkaline content which could improve human health; it is gluten-free, rich in micronutrients that play certain roles in the body immune system. Millets containing dietary is reported to act as antioxidants which can prevent deterioration of human health, also noted for some ailment prevention, like lowering blood pressure, risk of heart disease, prevention of cancer, cardiovascular diseases, diabetes, decreasing of tumor.

Keywords: Cereal; Millet; Sorghum; Soybean; Complementary Food

Introduction

Infant deaths accounted for Ten per cent of new-born deaths in the world last year occurred in Nigeria, a new report by the United Nations Children Fund, UNICEF, has revealed. World Health Organization [1,2]: confirmed that five countries accounted for half of all new-born deaths last year, with Nigeria third in the list. According to the report, India is ranked (24 per cent), Pakistan (10 per cent), Nigeria (9 per cent), the Democratic Republic of the Congo (4 per cent) and Ethiopia (3 per cent) [3]. Alarming report, new-born deaths occurred in two regions: Southern Asia (39%) and sub-Saharan Africa (38%) respectively. The report showed that 15,000 children died globally before their fifth birthday in 2016, with 46 per cent of the deaths (7,000) occurring in the first 28 days of life [1,2]. In a new study of World Health Organization entitled: Levels and Trends in Child Mortality 2017 (WHO) 2018. The study revealed that although the number of children dying before the age of five is reduced to 5.6 million in 2016 compared to nearly 9.9 million in 2000 - the proportion of under-five deaths in the new-born period has hiked up to 41 per cent to 46 per cent during the same period [1]. Similar report by

UNICEF also said though the lives of 50 million children under-five have been saved since 2000 through increased level of commitment by governments and development partners to tackle preventable child deaths, more still needs to be done to stop babies from dying the day they are born, or days after their birth. Joint report released by UNICEF, the World Health Organization, the World Bank and the Population Division of UNDESA, which make up the Inter-agency Group for Child Mortality Estimation (IGME), speculated that at current trends, 60 million children, will die before their fifth birthday between 2017 and 2030, half of them new-borns [1,2,4]. The Nigerian Minister of Health confirmed the high mortality rate of under-five in the country as frustrating and unacceptable and significant effort are made in reducing the rate of new-born deaths in the country as it has declined from 201/1000 live births to 128/1000 live births in 2013 [3]. The best measure of success for Universal Health Coverage is that every mother should not only be able to access health care easily, but that it should be quality, affordable care that will ensure a healthy and productive life for her children and family [3]. Preventable child deaths can be achieved by improving access to skilled health-professionals during pregnancy and at the time of birth; lifesaving interventions, such as immunization, breastfeeding and inexpensive medicines; and increasing access to water and sanitation giving adequate and balance dietary [3]. In most part of Africa, cereals supplies are amount to 80% of the energy requirements. Major cereals cultivate in Nigeria include rice, sorghum, maize, and pear millet; they are rich in carbon-hydrate but comparatively low in protein and naturally deficient in calcium and vitamin [5,6]. However many senero are attribute to infant mortality such as poor distribution of food in the family, taken poor protein, seasonal or severe shortage of food, poor supply of legumes, taboos and cultural practice, poor infant feeding practice (breast) and poverty [7-11]. Hence, we have embarked on research to formulate and determine of Nutritional composition of Cereal based complementary food as a succor to reduce child and maternal mortality rate in Nigeria.

Materials and Method

Some of the cereals used as source of energy in this experiment include maize, sorghum and millet. Source of protein include soybean, milk based commercial food were purchased from urban market in Ile- Ife, Nigeria.

Chemical analysis

The chemical analysis included Protein (nitrogen x 6.25), moisture, fat, crude fibre and carbohydrate of the ingredients, formulated diets and the nitrogen contents of the internal organ were determined according to AOAC (2005). Energy value was determined using Combustion calorimeter, model e2K [12-14].

Experimental animal procedure

Fifty white Wister albinos were procured from animal breeding centre; they were ranged from 46.55- 47.68g. They were four weeks old, respectively. The experiment animals were weighed randomly selected and distributed into five groups of ten per group and was housed in a metabolic cage [15,16]. They were fed on animal feeds (finisher) for seven days to acclimatize them to the new environment. The experimental animals were again reweighed and distributed into five groups of ten per group. 10g of supplements dietary were supplied daily. Remnants of the dietary samples were carefully recorded and the weights were noted. Weight gain/loss of the experimental animals was taken every three days and graphically sketched as in figure 1. At the end of the experiment, which was twenty-eight days, the experimental animals were anesthetized sacrificed. The organs collected from the animal including spleen, heart lungs kidney liver and small intestine were fixed immediately in 10% formyl saline for further experiment such as Nitrogen retention [7-11,15-20].

The food efficiency ratio (FER), Protein efficiency ratio (PER), the net protein retention (NPR) and protein retention efficiency (PRE) were calculated using the formula given below.

$$\text{The food efficiency ratio (FER)} = \frac{\text{Gain in body weight (g)}}{\text{Food intake (g)}}$$

$$\text{Protein efficiency ratio (PER)} = \frac{\text{Weight gain of test animal (g)}}{\text{Protein consumed by the test animal (g)}}$$

$$\text{Net protein retention (NPR)} = \frac{\text{Weight gain of test animal (g)} + \text{Average Weight Loss of Animal}}{\text{Protein consumed by the test animal (g)}}$$

$$\text{Protein retention efficiency (PRE)} = \text{NPR} \times 16$$

$$\text{Feed Conversion ratio was determined by} = \frac{\text{Feed Consumed (g)}}{\text{Gain in body weight (g)}}$$

Ethical consideration

Fifty white albino rats were approved for the experiment by the Animal Ethical Welfare Review Committee of the Obafemi Awolowo University, Osun State, Ile-Ife, Nigeria.

Results and Discussion

Sorghum 70% Soybean 30% (SS), Millet 70% Soybean 30% (MS), Sorghum 35% Millet 35%, Soybean 30% (SMS); Basal Dietary (BD) and Commercial Food (CF).

Figure 1 represented experimental Animal growth response for 28 days. The growth responses by experimental Animal were progressive by weight gained for dietary CF is 42.2g, MS is 32.2g, SMS is 19.22g, SS is 16.36g and but BD is declining by 3.2g. This shows that dietary experimental Animal growth contain essential nutrient such as complete amino acid profile [7-11,17,18,21-23].




Figure 1: Animal experimental growth for 28 days.

Table 1 showcased the chemical composition of the Formulated Dietary samples such as protein, it ranged from 15.34 - 16.01% moisture ranged from 12.5 - 5.68%, fat ranged from 0.50 - 10%, ash ranged from 11.45 - 2.30%, crude fibre ranged from 0.76 - 2.30%, dry matter ranged from 94.32 - 97.50% and Energy ranged from 364 - 413 Kcal%. The dietary formulation met the recommended daily intake and support animal growth fed with formulated sample dietary but BD could not support growth for lack of essential protein as earlier reported for lack of essential amino acid such as lysine and tryptophan hence could not support growth [7-11,21,24].

Dietary	Protein%	Moisture%	Fat%	Ash%	Crude Fibre%	CHO%	Dry matter%	Kcal%
SS	15.87 ± 01	5.53 ± 00	2.4 ± 02	1.45 ± 03	1.50 ± 01	73.25 ± 00	94.47 ± 02	383 ± 01
MS	15.30 ± 01	4.86 ± 01	0.35 ± 01	2.36 ± 02	0.80 ± 01	76.33 ± 03	95.14 ± 03	398 ± 01
SMS	15.34 ± 00	3.50 ± 03	0.56 ± 02	1.80 ± 01	0.78 ± 02	78.02 ± 02	96.50 ± 02	378 ± 02
BD	-	5.68 ± 01	0.50 ± 01	2.10 ± 00	0.76 ± 01	90.96 ± 01	94.32 ± 01	364 ± 03
CF	16.01 ± 02	2.50 ± 03	10 ± 01	4.30 ± 00	2.30 ± 00	64.89 ± 00	97.50 ± 00	413 ± 01

Table 1: Chemical composition of the formulated dietary samples. Mean ± SD values of five replicates (p ≥ 0.05); Sorghum 70% Soybean 30% (SS), Millet 70% Soybean 30% (MS), Sorghum 35% Millet 35%, Soybean 30% (SMS); Basal Dietary (BD) and Commercial Food (CF).

Table 2 is a portrait of the nitrogen retention in various tissues of the internal organ of the experimental animal liver, kidney and muscle. They are ranged from 18.34 - 72.04, 18.36 - 72.08 and 64.66 - 72.06 respectively. Commercial Food (CF) promoted the highest growth rate, followed by millet containing dietary such as Millet 70% Soybean 30% (MS), Sorghum 35% Millet 35%, Soybean 30% (SMS) also promoted the growth response similar to commercial diet. This is because Millet grain reported contains alkaline content it is gluten-free, rich and micronutrients that play certain roles in the body immune system. Millets is also reported to be antioxidants which can prevent deterioration of human health [23,25]. Millets is noted for some ailment prevention like lowering blood pressure, risk of heart disease, prevention of cancer, cardiovascular diseases, diabetes, decreasing of tumor. Alkaline based diet is often recommended to achieve optimal health. In developing countries, cereal-based foods have low bioavailability of minerals like iron, zinc initiate critical problem for infants and young children. Food processing techniques are used to enhance nutritional quality, improve the digestibility and bioavailability of food nutrients with reducing anti-nutrients [21,25,26].

Dietary	Kidney mg/g	Liver mg/g	Muscle mg/g
SS	65.30 ± 02	66.56 ± 01	64.66 ± 00
MS	70.66 ± 01	70.64 ± 02	70.68 ± 03
SMS	68.30 ± 02	64.56 ± 03	65.66 ± 04
BD	18.34 ± 01	18.36 ± 02	18.30 ± 01
CF	72.04 ± 01	72.08 ± 04	72.06 ± 03

Table 2: The nitrogen retention in various tissues of the internal organ. Mean ± SD values of five replicates (p ≥ 0.05); Sorghum 70% Soybean 30% (SS), Millet 70% Soybean 30% (MS), Sorghum 35% Millet 35%, Soybean 30% (SMS); Basal Dietary (BD) and Commercial Food (CF).

Table 4 reflected the Bioassay of the experimental animal Bioassay BV 75 - 66% NPU 64 - 72% PER 1.6 - 42 FER 1.6 - 4.2 and NPR 1.7 - 4.3, BD lacked biological value hence could not promote growth but Commercial food (CF) had the highest value, a bit lower than of white egg. It was followed by millet based dietary (MS) and SS and SMS [7-11,15-1821,27-30].

Dietary	Liver(g)	Kidney(g)	Body weight(g)	Gained(g) /loss
SS	2.80 ± 01	0.63 ± 02	70.86 ± 01	16.36 ± 02
MS	2.86 ± 00	0.73 ± 03	84.36 ± 01	32.2 ± 03
SMS	3.43 ± 03	0.66 ± 00	79.63 ± 02	19.22 ± 04
BD	2.8 ± 02	0.45 ± 01	58.25 ± 02	3.2 ± 02
CF	5.1 ± 01	0.80 ± 01	104.06 ± 00	42.2 ± 01

Table 3: Nitrogen retention of the experimental animal.

Mean ± SD values of five replicates ($p \geq 0.05$); Sorghum 70% Soybean 30% (SS), Millet 70% Soybean 30%(MS), Sorghum 35% Millet 35%, Soybean 30% (SMS), Basal Dietary (BD) and Commercial Food (CF).

Diet	BV %	NPU%	PER	NPR	Gained/loss
SS	66 ± 01	64 ± 0.2	1.6 ± 0.2	1.7±0.3	16.36 ± 0.3
MS	73 ± 00	70 ± .1	3.2 ±	3.3 ±	32.2 ±
SMS	66 ± 00	65 ± .01	1.9 ± .02	2.0 ± 02	19.22 ± 01
BD	-	-	-	-	-3.2 ± 01
CF	75 ± 0	72 ± .0	4.2 ± 01	4.3 ± 02	42.2 ± 0

Table 4: Bioassay of the experimental animal Bioassay BV % NPU% PER and NPR.

Mean ± SD values of five replicates ($p \geq 0.05$); sorghum 70% soybean 30% (ss), millet 70% soybean 30% (ms), sorghum 35% millet 35%, soybean 30% (sms), basal dietary (bd) and commercial food (cf).

Conclusion

In conclusion cereal based complementary are produced from staple food are available for baby and infant in most part of the country. Millet grain containing diets promoted the experimental animal growth response similar to commercial diet. It contains alkaline content which could improve human health; it is gluten-free, rich in micronutrients that play certain roles in the body immune system. Millets containing diet is reported to act as antioxidants which can prevent deterioration of human health and also noted for some ailments prevention, like lowering blood pressure, risk of heart disease, prevention of cancer, cardiovascular diseases, diabetes, decreasing of tumor.

Acknowledgments

The authors are very grateful to staff of Department of Food Science and Technology, Joseph Ayodele Babalola University and Department of Food Science and Technology, Obafemi Awolowo University, Ile-Ife, Nigeria for their collaboration in this work.

Bibliography

1. World Health Statistics series (WHO). 36 health-related Sustainable Development Goal (SDG) indicators (2018).
2. UNICEF. Levels and Trends in Child Mortality Report (2017).
3. Vangurd newspaper (VNP, 2017) (2018).
4. World Bank Group (WBG). "Levels and trends in child mortality: report 2017". Washington, D.C (2017).
5. FAO. "Food and Agriculture Organization of the United Nations: Fisheries and Aquaculture Department". Rome, Italy The state of world Aquaculture and Fisheries (2007).

6. Foramfera. "Maize Production and Sales in Nigeria: The Opportunity". Foramfera (2013).
7. Ibiroinke SI and Adepeju AB. "Nutritional Assessment of Breakfast Foods Developed from Animal Polypeptide, Crayfish (*Euastacus Spp*) and Maize (*Zea Mays*)."
American Journal of Food and Nutrition 4.5 (2016a): 131-134.
8. Ibiroinke SI., et al. "Histopathological And In-Vivo Study of Moringa Oleifera Seed".
Annals Food Science and Technology 17.1 (2016b): 102-108.
9. Ibiroinke S I., et al. "Comparative Study of Condiment Vegetable Basil Leaf (*Ocimum gratissimum*) and Bitter Leaf (*Vernonia Amygdalina*)."
American Journal of Food and Nutrition 5.3 (2017a): 95-98.
10. Ibiroinke SI., et al. "Haematological and In-Vivo Study of Moringa Oleifera Seed".
MOJ Food Process Technol 4.6 (2017b): 00109.
11. Ibiroinke S I., et al. "Comparative Study of Condiment Vegetable Basil Leaf (*Ocimum gratissimum*) and Bitter Leaf (*Vernonia Amygdalina*)."
American Journal of Food and Nutrition 5.3 (2017c): 95-98.
12. Oladeji BS., et al. "Physico-Chemical And Nutritional Evaluation of Co-Processed Fermented Yellow Maize Ogi (An Infant Diet) And Carrot Blends".
Annals Food Science and Technology 15.1 (2014): 82-91.
13. Oladeji BS., et al. "Comparative analysis of physico-chemical properties and amino acids profile of three tropical maize hybrid cultivars in Nigeria Emerald".
Nutrition and Food Science 46.5 (2016): 695-705.
14. AOAC. Official of Analysis Methods of Analysis of the Association of Official Analytical Chemists, 18th Edition (2005).
15. Ibiroinke SI., et al. "Nutritional Evaluation of Complementary Food Developed from Plant and Animal Protein Sources".
Emerald Nutrition and Food Science 42.2 (2012a): 111-120.
16. Ibiroinke SI. "Formulation of Infant Weaning Foods from Vegetable Proteins and Cereal".
American Journal of Food Technology 9.2 (2014a):104-110.
17. Ibiroinke SI and Ige MM. "Nutritional Evaluation and Chemical Analysis of Two Commercial Infant Foods in South-Western, Nigeria".
International Journal of Nutrition and Food Sciences 4.5 (2015a): 590-593.
18. Ibiroinke SI and Fawale O. "Chemical Analysis and Animal Bioassay of *Hildegardia barteri* (Kariya Seed) SoyBean and Maize".
Annals Food Science and Technology 16.1 (2015b): 245-250.
19. Kuku A Alabi and MO Ibiroinke SI. "Effect of processing on the lectin and trypsin inhibitor content of *Plukenetia conophora* seeds as it affects growth performance and nutrient metabolism in rat".
African Journal of Food Science 7.9 (2013): 306-316.
20. Kuku A., et al. "Processing of Fluted Pumpkin Seeds, *Telfairia Occidentalis* (Hook F) As It Affects Growth Performance and Nutrient Metabolism in Rats".
African Journal of Food and Agriculture Nutrition and Development 14.5 (2014): 1992-2014.
21. Titani K., et al. "Amino acid sequence of crayfish (*Astacus fluviatilis*) trypsin If".
Biochemistry 22.6 (1983):1459-1465.
22. Sarita and Ekta Singh. "Potential of Millets: Nutrients Composition and Health Benefits".
Journal of Scientific and Innovative Research 5.2 (2016): 46-50.
23. Shahidi F and Chandrasekara A. "Millets grain phenolics and their role in disease risk reduction and health promotion: A review".
Journal of Functional Foods 5.2 (2013): 570-581.
24. Food and Agriculture Organization (FAO). "Crops statistics – concepts, definitions and classifications?" (2010).

25. Lei V., *et al.* "Spontaneously Fermented Millet Product as a Natural Probiotic Treatment for Diarrhea in Young Children: An Intervention Study in Northern Ghana". *International Journal of Food Microbiology* 110.3 (2006): 246–253.
26. Oloyede FM., *et al.* "Antioxidant Activities and Food value of Five Underutilized Green Leave Vegetables in South Western Nigeria". *The Nigerian Journal of Nutritional Science* 32 (2011): 13-18.
27. Delgado C., *et al.* "Fish to 2020. Supply and demand in changing global market". International Food Policy Research Institute. Washington D.C (2003).
28. Ibiroinke SI and Alakija O. "Production of Soy Cheese from Vegetable Protein Using Different Coagulants". *Journal of Nutrition and Health Sciences* 5.1 (2018a): 107.
29. Ibiroinke SI., *et al.* "Nutritional Evaluation and Comparison Study of Seafoods, Fish and Crayfish Dietary". *MOJ Food Process Technology* 6.21 (2018b): 00109.
30. Ibiroinke SI., *et al.* "In vivo and haematological study of some selected vegetable condiments". *MOJ Food Process Technology* 6.3 (2018): 318–323.

Volume 5 Issue 8 August 2019

©All rights reserved by Adefisola Bola Adepeju and Samson Ishola Ibiroinke.