

The Association of Low Dietary Diversity Score with the Nutritional Status of Expectant Mother from Rural Ethiopia: A Community Based Cross-Sectional Study

Nigussie Assefa Kassaw¹, Kibret Ephram² and Jemal Haidar^{1*}

¹School of Public Health, College of Health Sciences, Ethiopia

²Former MPH Student, Ethiopia

***Corresponding Author:** Jemal Haidar, School of Public Health, College of Health Sciences, Ethiopia.

Received: February 06, 2021; **Published:** September 28, 2021

Abstract

Background: Maternal underweight is a challenging public health issues globally affecting women in most developing countries and consequently affect their newborns. Such information however is scanty in the study area and thus we examined the association of low dietary diversity score with the nutritional status of expectant mother from rural Ethiopia.

Methods: This study was based data extracted from a large data set gathered from three rural villages from September 2017 to June 2018 in a district of North-Shewa Zone of Amhara Regional State which is 220 km north of Addis Ababa, the capital city of Ethiopia. Data on socio demographic characteristics food groups consumed and MUAC measurement were taken for all expectant mothers by trained health extension workers in accordance with relevant ethical guidelines and regulations. The collected data were entered into Epi-data software version 3.02 and analyzed using SPSS version 20. Statistical significance was affirmed at a p-value ≤ 0.05 .

Result: Of the 400 consented participants, 380 of them had adequate information and thus analyzed for their variety of food groups consumption and their MUAC measurement outcomes. The magnitude of underweight was 18.2% and the mean (\pm SD) dietary diversity score was 4.45 (± 1.32). interestingly, the relationship between poor dietary diversity score and underweight of the participants was significant (AOR = 2.4; 95% CI = 1.05 to 5.3).

Conclusion: Our study revealed a significant relationship between poor dietary diversity score and underweight of expectant mothers. it is thus, essential to improve the dietary diversity and nutritional status of pregnant women through strengthening the ongoing productive safety net program to earn more income that would enable mothers to purchase variety of foods and ultimately break the intergeneration cycle of malnutrition.

Keywords: Children; Mothers; Vitamin A Deficiency; Night Blindness; Prophylaxis; Sudan

Introduction

Maternal underweight is a challenging public health issues globally affecting women in most developing countries including Ethiopia and responsible for the death of mothers and children [1-4]. Because of the grave consequences of malnutrition women are advised to take more diversified food to get adequate energy and multiple nutrients to satisfy their demand as well as their offspring. Mothers in most developing countries however, do not adhere to the nutritional advices provided to them during their antenatal follow up due to

several factors that include limited access to and consumption of micronutrient rich foods, frequent illnesses, teenage pregnancy with narrow birth intervals and high fertility rates [1,6].

The solution requires effective multisectoral interventions that include health, social welfare, education, water, sanitation, in addition to consumption of balanced diet which contributes to the wellbeing of the mother as well as her offspring [7]. In line with this, the government of Ethiopia has developed a revised national nutrition program (NNP) in 2016 to address the burden of malnutrition in pregnant and lactating women [8].

Despite the implementation of the revised NNP initiative, maternal underweight remained a pressing problem according to the 2016 Ethiopian demographic health survey which documented 22.4% of women in reproductive ages (15 - 49 years) were undernourished [9].

Other than the aforementioned documented report, several pocket studies in the country also revealed underweight to prevail and ranged from 9.2% to 31.8% [10-14].

Other recent studies done on pregnant women indicated that poor dietary scores in the same age and sex groups are documented to be another important problem that have a strong positive relationship between appropriate maternal nutritional status and birth outcomes and ranged from 24.2% - 74.6% [15-17].

Nonetheless, such information for expectant mothers from rural north Shewa is scanty and therefore, examined the association of low dietary diversity score with the nutritional status of pregnant women based on a large data set collected in a rural community to generate evidence that can be used to design appropriate interventions for the villages as well as for the region and the country.

Methods

This study was extracted from a large data set gathered in a Community based cross-sectional study conducted among expectant mothers from September 2017 to June 2018 in a rural district of North-Shewa Zone of Amhara Regional State which is 220 km north of Addis Ababa, the capital city of Ethiopia [Master Thesis, 2018].

A total of 400 expectant mothers from five villages were randomly selected in accordance with relevant ethical guidelines and regulations in the initial mother document and allocated proportionally to each of the villages based on the constituted sampling frame obtained from health posts.

Data on socio demographic characteristics food groups consumed (Nine food groups as displayed in table 1) and MUAC measurement were taken for all expectant mothers by trained health extension workers. Maternal MUAC was measured using non stretchable tape (MUAC) and recorded as normal or abnormal when MUAC > 22 cm and < 22 cm, respectively. Dietary diversity scores (DDS) was formed by summing up the number of food groups consumed over a 24-hour period by an individual and labeled as high and low DDS when the food groups consumed were more than five and below five, respectively as recommended by FAO and FHI 2016 [18]. The collected data were entered into Epi-data software version 3.02 and analyzed using SPSS version 20.

Result

Of the 400 recruited mothers, only 380 of them who had complete information were included in the analysis and the rest were excluded.

Consumption of the modified dietary diversity score (MDDS)

Grain, white roots and tubers were amongst the commonly consumed grains by all expectant mothers predominantly in the form of breads, pasta, macaroni, porridge, rice. The second most commonly consumed meals as their integral part of the main dish for the vast

majority (98.7%) was vegetables. Consumption of pulses and other vitamin A-rich fruits/vegetables was also reported in 66.3% and 55.3% of them, respectively. Milk and milk products (dairy), and other dark green leafy vegetables were consumed in 46.6% and 29.7%, respectively. Meat, poultry and fish were consumed in 27.6%. the details of food groups are displayed in table 1.

Food groups	Frequency N = 380	Percentage (%)
1. Grains, white roots and tubers, and plantains	380	100
2. Other vegetables	375	98.7
3. pulses (beans, peas and lentils)	252	66.3
4. Other vitamin A-rich fruits and vegetables	210	55.3
5. Dairy	177	46.6
6. Dark green leafy vegetables	113	29.7
7. Meat, poultry and fish	105	27.6
8. Egg	62	16.3
9. Other fruits	17	4.5

Table 1: Consumption of the MDDS of the respondents by nine food groups.

As shown in table 2, the proportion of mothers who consumed more than five food groups (high dietary diversity score) was nearly 43.0% while those who consumed five or less (low dietary diversity score) were close to 57.0%. The mean DDS was 4.45 ± 1.32.

Dietary diversity scores	N (%)
G Consumption of 5 or less food groups (low DDS)	217 (57.1%)
Consumption of more than 5 food groups (high DDS)	163 (42.9%)
Mean (SD)	4.45 (1.32)

Table 2: Dietary diversity score of the expectant mothers.

Underweight level determined by MUAC and its association with dietary diversity score

As displayed in table 3, a large number (81.8%) of expectant mothers had optimum nutritional status while less than a quarter (18.2%) were underweight based on the suggested cut off point for MUAC. The proportion of underweight mothers was higher in the age group of 15 to 24 years than those above 25 years and the difference noted was statistically significant in the bivariate analysis (COR = 0.56; 95%CI = 0.03 to 0.95). Nevertheless, this association when further analyzed, its significance was lost in the multivariate regression analysis.

Characters	Normal	Underweight	COR (95% CI)	AOR (95% CI)
Age (in years)				
15 - 25	136 (77.3%)	40 (22.7%)	0.56 (.033 - .955)	0.18 (.22 - 1.5)
Greater than 25	175 (85.8%)	29 (14.2%)	1	1
Total	311 (81.8%)	69 (18.2%)		
Dietary diversity score				
≥ 5	131 (80.4%)	32 (19.6%)	0.841 (.498 - 1.421)	2.4 (1.056 - 5.3)*
< 5	180 (82.9%)	37 (17.1%)	1	1
Total	311 (81.8%)	69 (18.2%)		

Table 3: Association of dietary diversity scores and nutritional status.

COR: Crude Odd Ratio; AOR: Adjusted Odd Ratio.

As expected on the other hand, DDS was significantly associated with the nutritional status of the expectant mothers in the multivariate analysis (AOR = .4; 95% CI = 1.05 to 5.3).

Discussions

The association of low dietary diversity score with the nutritional status of expectant mother from rural Ethiopia was examined and showed the overall proportion of underweight to be less than a quarter and the majorities to have low DDS. Nonetheless, when the proportion of underweight is disaggregated against age, the most affected age group was those in the age group between 15 and 25 years. In the same breath, the proportion of underweight was significantly higher among mothers with low DDS indicating low dietary diversity is a significant factor associated with the nutritional status of mothers.

The magnitude of underweight observed in the present study is concordant with Northwest Ethiopia [19], Sri Lanka [20] and Kenya [21] which documented 14.4%, (15%) and (19.3%) of underweight among pregnant mothers, respectively.

On the other hand, when compared with the Indian (61.5%) and South West Ethiopia (44.9%), our finding is expressively lower than the finding documented for India [22] and south west Ethiopia [27].

The difference might be due to that the study in India exclusively included participants from lower social classes. Further, the discrepancies might be attributed to the difference in stage of pregnancy, nutritional practice, a seasonal variation on food consumption and the use of different cut off points for mid-upper arm-circumference measurement in different studies.

The proportion of mothers with high DDS in our study is better than that of Pakistani report (80.4 vs 5.0) which documented only 5% of mothers to have high DDS [24].

In contrast, when compared with the Kenyan study our finding is lower (80.4 vs 92.4) [20]. Such variations are expected due to the differences in the sociodemographic status and to some extent to the cultural reasons of the participants. The mean DDS (\pm SD) observed in our study is closer to the findings of pregnant women reported mothers in Southwest Ethiopia (4.6 vs 4.3) [26], Bangladesh (4.6 vs 4.38) [27] and higher than the study finding from central Ethiopia (4.6 vs 3.5) [28]. On the other hand, compared with the Kenyan which reported a mean DDS of 6.8 [29] and South Ethiopian mean DDS of 7.0 [30], the present figure is however lower than the above-mentioned findings possibly due to their differences in socio-socioeconomic status, and cultural/seasonal variations.

As in many developing countries, this study showed that plant-based foods are the major sources of both macro and micronutrient for the vast majority of the mothers and were consumed in the form of breads, pasta, macaroni, porridge, rice followed by vegetables. While consumption of animal-based proteins which are good sources of nutrients is markedly low and consequently increased the risk of nutritional deficiencies. Similar findings were reported Kenya and southwest Ethiopia and Eastern Ethiopia [12, 26,28, 29,31-34].

In this study, less than a quarter of the expectant mothers were underweight based on the suggested cut off point of MUAC measurement of less than or equal to 22 cm and most of the underweight cases were among the younger age group and our finding is concordant with some studies conducted in the country and elsewhere. For instance, in Northwest Ethiopia, Sri Lanka and Kenya [19-23,26], the magnitude of underweight among same groups was reported to be 14.4%, 15% and 19.3%, respectively. When compared with the Indian study [22], our finding is markedly lower (18.7 vs 61.5) and this difference is probably attributed to cultural and economic factors as most of the cases in India were from the lower social classes.

As expected, the odds of underweight were 4 times higher among mothers with poor DDS than their referent groups and our finding is concordant with some previous pocket studies conducted locally [26,33]. It is eminent that when women consume less food groups, particularly low consumption of animal-based proteins which are good sources of nutrients, there would be an increased risk for malnutrition and similar finding is reported in west Ethiopia [26,32].

The strength of this study is that we used rural community that had no data collection fatigue and validated tool. Nonetheless, the study had some limitations emanated from the nature of study design which might led to recall bias.

Conclusion

In conclusion, our study revealed poor dietary diversity practices to be significantly associated with the observed level of maternal underweight. It is thus, essential for all development sectors to improve the dietary diversity and nutritional status of pregnant women through strengthening the current productive safety net program to earn more income that would enable mothers to purchase variety of foods and ultimately break the intergeneration cycle of malnutrition. Other than this, mothers need to be counselled on the importance of the nutritional value of different food groups and its use.

Bibliography

1. Zulfiqar AB., *et al.* "Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost?" *The Lancet* (2013): 40-65.
2. Robert EB., *et al.* "Maternal and child undernutrition: global and regional exposures and health consequences". *The Lancet* (2008): 5-22.
3. Alemayehu MS and Tesema EM. "Dietary Practice and Associated Factors among Pregnant Women in Gondar Town North West". *Ethiopia International Journal of Nutrition and Food Sciences* 4.6 (2015): 707-712.
4. Dereje Tsegaye., *et al.* "Factors Associated with Dietary Practice and Nutritional Status of Pregnant Women in Rural Communities of Illu Aba Bor Zone, Southwest Ethiopia". *Nutrition and Dietary Supplements* 12 (2020): 103-112.
5. Shiferaw A and Husein G. "Acute under nutrition and associated factors among pregnant women in Gumay District, Jimma Zone, South West Ethiopia". *Journal of Women's Health Care* 8.2 (2019): 459.
6. Ethiopian Public Health Institute. Ethiopian national food consumption survey. EPHI (2013): 1-57.
7. Fouda LM., *et al.* "Nutritional Awareness of Women during Pregnancy". *Journal of American Science* 8.7 (2012).
8. FDRE. Government of Ethiopia, National Nutrition Program 2016-2020.
9. CSA and ICF. Ethiopia Demographic and Health Survey. Addis Ababa, Ethiopia, and Rockville". Maryland, USA (2016).
10. Dadi AF., *et al.* "Undernutrition and its associated factors among pregnant mothers in Gondar town, Northwest Ethiopia". *PLoS One* 14.4 (2019): 1-11.
11. Nigatu M., *et al.* "Household food insecurity, low dietary diversity, and early marriage were predictors for undernutrition among pregnant women residing in Gambella, Ethiopia". *Hindawi Advances in Public Health* (2018): 1-10.
12. Kuche D., *et al.* "Nutritional status and associated factors among pregnant women in Wondo Genet District, Southern Ethiopia". *Journal Food Science and Engineering* 5 (2015): 85-94.
13. Zema DT. "Factors associated with dietary practice and nutritional status of pregnant women in Dessie town, northeastern Ethiopia: a community-based cross-sectional study". *BMC Pregnancy Childbirth* 19.1 (2019): 517.
14. Mariyam AF and Dibaba B. "Journal of nutritional disorders and epidemiology of malnutrition among pregnant women and associated factors in Central Refit Valley of Ethiopia, 2016". *Journal of Nutritional Disorders and Therapy* 8.1 (2018): 1-8.
15. Jemal K and Awol M. "Minimum dietary diversity score and associated factors among pregnant women at Alamata General Hospital, Raya Azebo Zone, Tigray Region, Ethiopia". *Journal of Nutrition and Metabolism* (2019): 1-6.

Citation: Jemal Haidar., *et al.* "The Association of Low Dietary Diversity Score with the Nutritional Status of Expectant Mother from Rural Ethiopia: A Community Based Cross-Sectional Study". *EC Nutrition* 16.10 (2021): 55-60.

16. Nana A and Zema T. "Dietary practices and associated factors during pregnancy in northwestern Ethiopia". *BMC Pregnancy Childbirth* 18.1 (2018): 183.
17. Desta M., et al. "Dietary diversity and associated factors among pregnant women attending antenatal clinic in Shashemane, Oromia, Central Ethiopia: a cross-sectional study". *Journal of Nutrition and Metabolism* (2019): 7.
18. FAO and FHI. "Minimum Dietary Diversity for Women a Guide to Measurement (2016): 360.
19. Desyibelew HD., et al. "Burden and determinants of malnutrition among pregnant women in Africa: a systematic review and meta-analysis". *Plos One* 14.9 (2019): 1-19.
20. Adikari AMNT., et al. "Assessment of nutritional status of pregnant women in a rural area in Sri Lanka". *Tropical Agricultural Research* 27.2 (2016): 203-211.
21. Willy K., et al. "Dietary diversity, nutrient intake and nutritional status among pregnant". *International Journal of Health Sciences* 6 (2016): 378-385.
22. Nisal KM. "Women belonging to low income group". *Journal of Scientific* 5.8 (2015): 580-582.
23. Ali F., et al. "Assessment of dietary diversity and nutritional status of pregnant women in islamabad, pakistan". *Journal of Ayub Medical College Abbottabad* 26.4 (2014).
24. Kiboi W., et al. "Dietary diversity, nutrient intake and nutritional status among pregnant women in Laikipia County, Kenya". *International Journal of Health Sciences and Research* (2016): 378-379.
25. Dereje T., et al. "Factors Associated with Dietary Practice and Nutritional Status of Pregnant Women in Rural Communities of Illu Aba Bor Zone, Southwest Ethiopia". *Nutrition and Dietary Supplements* 12 (2020): 103-112.
26. Shamim AA., et al. "Pregnant women diet quality and its sociodemographic determinants in Southwestern Bangladesh". *Food and Nutrition Bulletin* 37.1 (2016): 14-26.
27. Desta M., et al. "Dietary diversity and associated factors among pregnant women attending antenatal clinic in Shashemane, Oromia, Central Ethiopia: a cross-sectional study". *Journal of Nutrition and Metabolism* (2019): 7.
28. Willy K., et al. "Dietary diversity, nutrient intake and nutritional status among pregnant". *International Journal of Health Sciences* 6 (2016): 378-385.
29. Romedan Delil DT. "Dietary diversity and its association with anemia among pregnant women attending public health facilities in South Ethiopia". *Ethiopian Journal of Health Sciences* 28.5 (2018): 625-634.
30. Hailu S and Woldemichael B. "Dietary diversity and associated factors among pregnant women attending antenatal care at public health facilities in Bale Zone, Southeast Ethiopia". *Nutrition Dietary Supplements* 11 (2019): 1.
31. Moges A., et al. "Under nutrition and Associated Factors among Adolescent Pregnant Women in Afdem district, Ethiopian Somali Region, Eastern Ethiopia". Haramaya University (2017).
32. Shenka A., et al. "Dietary Diversity and Nutritional Status of Pregnant Women Attending Public Hospitals in Dire Dawa City Administration, Eastern Ethiopia". *East African Journal of Health Biomedical Sciences* 2.1 (2018): 10-17.

Volume 16 Issue 10 October 2021

©All rights reserved by Jemal Haidar., et al.